



For eczema, itching, and  
frostbite

Eczema, dermatitis treatment medicine

Product Name: Rinderon Vs Ointment

# Rinderon Vs Ointment

Class 2 Type Drug



This drug is a skin disease treatment, so do not use it under makeup or after shaving.  
Using large amounts or for a long period of time may cause side effects such as thinning of the skin and dilation of blood vessels in the skin. Take special care with the skin on the face, as it is thin.  
In addition, do not continue using the drug indiscriminately after symptoms have improved.



Please read  
carefully.



## PRECAUTIONS



When not to use this product

(If do not follow these instructions, current symptoms may worsen or adverse reactions are more likely to occur.)

- This product should not be used in the following persons.**  
Patients who have had an allergic symptom to this drug or its ingredients
- The medicine should not be used on the following parts:**
  - Varicella (chickenpox), athlete's foot/tinea or purulent lesions
  - Eyes and periorbital area
- Do not take extensively on the face.**
- Do not take this medicine for a long time.**



## Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.**  
(1) Patients undergoing medical treatment from a doctor (2) Pregnant women or women suspected of being pregnant (3) Patients who have experienced allergic symptoms associated with drugs, etc. (4) Those with extensive lesions (5) Those with severe moistening or erosion
- If the following symptoms are observed after using this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.**
- When symptoms do not improve even after using this drug medicine 5 to 6 times, stop using the drug and consult a physician, pharmacist or registered salesperson, being sure to take this document with you.**

| Affected area          | Symptom  |
|------------------------|--|
| Skin                   | Rash/redness, itching  |
| Skin (affected lesion) | Dermatomycosis such as athlete's foot/tinea, acne, purulent symptoms, persistent irritation, whitening |



## Indications

Eczema, dermatitis, miliaria, irritated skin, itching, chilblain, frostbite, urticaria



## Dosage and Administration

Apply an appropriate amount to the affected lesion once or several times a day.

<Precautions for dosage and Administration>

- Strictly follow the prescribed dosage and administration.
- When allowing children to use the medicine, a guardian must be present to watch and tell such them how to take it.
- Attention should be paid so that the medicine will not come into contact with eyes. If the medicine comes into contact with eyes, immediately rinse them with water or lukewarm water. If the symptoms are severe, seek an ophthalmologist for medical treatment.
- This medicine is for external use only.
- Do not cover the area with something that does not breathe, such as plastic wrap. When using on areas that come into contact with a diaper, do not use with tight-fitting diapers or sealed pants such as vinyl.
- Do not use under makeup or after shaving.



## Ingredient and amounts

Rinderon Vs Ointment is a white to pale yellow ointment, containing the following amount of ingredient per 1 gram.

| Ingredient             | Per 1g | Mode of action   |
|------------------------|--------|--|
| Betamethasone valerate | 1.2 mg | A corticosteroid to suppress inflammation and to relieve itching |

### Additives:

Liquid paraffin and white petrolatum



## Precautions for storage and handling

- Store in a cool, low humidity place away from direct sunlight.
- Keep out of reach of children.
- Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the product.)
- Do not use products after its expiration date.



## Contact Desk

For inquiries about this product, please contact the store where you purchased it or contact the following:

● Shionogi Healthcare Co., Ltd. "Drug Information Center"

Tel: Osaka 06-6209-6948, Tokyo 03-5406-8450

Reception hours: 9:00-17:00 (excluding Saturdays, Sundays and holidays)

For information on the "Relief System for Sufferers from Adverse Drug Reactions"

● Pharmaceuticals and Medical Devices Agency (PMDA)

[https://www.pmda.go.jp/kenkouhigai\\_camp/index.html](https://www.pmda.go.jp/kenkouhigai_camp/index.html)

Tel: 0120-149-931 (toll-free number)

# Rinderon Vs Ointment

**The key to treating skin problems is to heal quickly.**

**Topical steroids are effective for skin problems such as eczema and dermatitis.**

Skin problems such as dermatitis, eczema, rashes, and the associated itching are caused by inflammation of the skin. If the unpleasant itchy symptoms persist for a long time, and continue to scratch skin, which can worsen the symptoms and leave scars. The key to treating skin problems is to suppress inflammation early and alleviate symptoms as quickly as possible. Therefore, it is important to use topical steroids correctly.

Steroids are substances produced in the human body that have strong anti-inflammatory effects. Topical steroids, which are chemically synthesized steroids, are medications that reduce localized inflammation.

**It is effective against everyday skin problems and various types of dermatitis.**



※Do not use if the affected area is infected and oozing.


## Use for skin symptoms associated with suppuration

Dermatitis often causes severe itching, and continued scratching weakens the barrier function that protects the skin from infection, making it easier for bacteria to enter through the skin. This medication improves symptoms such as dermatitis and itching. Do not use on areas with suppuration.

| Condition    | Dermatitis without Suppuration (Infection) | Dermatitis with Suppuration (Infection) |
|--------------|--|---|
| Use or Don't | ○  | ×                                       |

## How to open the tube for the first time

Open the cap, turn it upside down, and insert the protrusion on the cap firmly into the opening of the tube to make a hole.



## Point

### Apply an appropriate amount

The amount that can be applied to an area the size of two palms is roughly the amount of medicine placed on an adult's index finger from the tip to the first joint (approximately 0.5g).



### Apply gently

Apply the ointment on the affected area and gently spread it over the entire affected area without rubbing it in. The basic rule is to apply it until the surface becomes sticky.



If symptoms do not improve or worsen after 5 to 6 days of use, stop using the medicine and consult a doctor, pharmacist, or registered seller.

## Side Effects

The frequency of side effects is low when topical steroids are used properly such as by the period of use and the amount applied. If used for a long period, the skin may become pale or thin. In addition, if used on areas infected with athlete's foot, the symptoms may worsen.

Does not aggravate skin problems  
Attention for daily life

- ✓ The itching worsens when body gets warm, so bathe in lukewarm water for a short time.
- ✓ Wash body gently with a soft towel or your hands.
- ✓ Avoid clothing and bedding that come into direct contact with the skin and are made from materials such as wool or synthetic fibers.
- ✓ Avoid stimulants such as alcohol and spicy foods.

