



For eczema, itching, and frostbite

Eczema, dermatitis treatment medicine Rinderon Vs Cream

Product Name: Rinderon Vs Cream

Class 2 Type Drug



This drug is a skin disease treatment, so do not use it under makeup or after shaving. Using large amounts or for a long period of time may cause side effects such as thinning of the skin and dilation of blood vessels in the skin. Take special care with the skin on the face, as it is thin. In addition, do not continue using the drug indiscriminately after symptoms have improved.



Please read carefully.



PRECAUTIONS



When not to use this product

(If do not follow these instructions, current symptoms may worsen or adverse reactions are more likely to occur.)

- This product should not be used in the following persons.**
Patients who have had an allergic symptom to this drug or its ingredients
- The medicine should not be used on the following parts:**
 - Varicella (chickenpox), athlete's foot/tinea or purulent lesions
 - Eyes and periorbital area
- Do not take extensively on the face.**
- Do not take this medicine for a long time.**



Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.**
(1) Patients undergoing medical treatment from a doctor (2) Pregnant women or women suspected of being pregnant (3) Patients who have experienced allergic symptoms associated with drugs, etc. (4) Those with extensive lesions (5) Those with severe moistening or erosion

- If the following symptoms are observed after using this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.**

Affected area	Symptom
Skin	Rash/redness, itching
Skin (affected lesion)	Dermatomycosis such as athlete's foot/tinea, acne, purulent symptoms, persistent irritation, whitening

- When symptoms do not improve even after using this drug medicine 5 to 6 times, stop using the drug and consult a physician, pharmacist or registered salesperson, being sure to take this document with you.**



Indications

Eczema, dermatitis, miliaria, irritated skin, itching, chilblain, frostbite, urticaria



Dosage and administration

Apply an appropriate amount to the affected lesion once or several times a day.

<Precautions for dosage and Administration>

- Strictly follow the prescribed dosage and administration.
- When allowing children to use the medicine, a guardian must be present to watch and tell such them how to take it.
- Attention should be paid so that the medicine will not come into contact with eyes. If the medicine comes into contact with eyes, immediately rinse them with water or lukewarm water. If the symptoms are severe, seek an ophthalmologist for medical treatment.
- This medicine is for external use only.
- Do not cover the area with something that does not breathe, such as plastic wrap. When using on areas that come into contact with a diaper, do not use with tight-fitting diapers or sealed pants such as vinyl.
- Do not use under makeup or after shaving.



Contact Desk

For inquiries about this product, please contact the store where you purchased it or contact the following:

● Shionogi Healthcare Co., Ltd. "Drug Information Center"
Tel: Osaka 06-6209-6948, Tokyo 03-5406-8450
Reception hours: 9:00-17:00 (excluding Saturdays, Sundays and holidays)



Ingredient and amounts

Rinderon Vs Cream is a white cream, containing the following amount of ingredient per 1 gram:

Ingredient	Per 1g	Mode of action
Betamethasone valerate	1.2 mg	A corticosteroid to suppress inflammation and to relieve itching

Additives:

White petrolatum, liquid paraffin, cetanol, polyoxyethylene stearyl ether, butyl parahydroxybenzoate, methyl parahydroxybenzoate, sodium dihydrogenphosphate, phosphoric acid, and sodium hydroxide



Precautions for storage and handling

- Store in a cool, low humidity place away from direct sunlight.
- Keep out of reach of children.
- Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the product.)
- Do not use product after its expiration date.



For information on the "Relief System for Sufferers from Adverse Drug Reactions"

● Pharmaceuticals and Medical Devices Agency (PMDA)
https://www.pmda.go.jp/kenkouhigai_camp/index.html
Tel: 0120-149-931 (toll-free number)

Rinderon Vs Cream

The key to treating skin problems is to heal quickly.

Topical steroids are effective for skin problems such as eczema and dermatitis.

Skin problems such as dermatitis, eczema, rashes, and the associated itching are caused by inflammation of the skin. If the unpleasant itchy symptoms persist for a long time, and continue to scratch skin, which can worsen the symptoms and leave scars. The key to treating skin problems is to suppress inflammation early and alleviate symptoms as quickly as possible. Therefore, it is important to use topical steroids correctly.

Steroids are substances produced in the human body that have strong anti-inflammatory effects. Topical steroids, which are chemically synthesized steroids, are medications that reduce localized inflammation.


It is effective against everyday skin problems and various types of dermatitis.



※Do not use if the affected area is infected and oozing.

Topical steroids are classified into 5 Levels

Topical steroids are classified into five levels based on the strength of their effect. Over-the-counter medications come in three levels: "weak," "mild," and "strong," and the betamethasone valerate contained in this product is classified as "strong."



No antibiotics included

Dermatitis often causes severe itching, and continued scratching weakens the barrier function that protects the skin from infection, making it easier for bacteria to enter through the skin. This medication improves symptoms such as dermatitis and itching. Do not use on areas with suppuration.

Condition	Dermatitis without Suppuration (Infection)	Dermatitis with Suppuration (Infection)
Use or Don't	○	×

Point

HOW TO USE

Apply an appropriate amount

If the amount of medicine is a little, it will not be effective. Apply the appropriate amount, as shown below, once or several times a day.

Usage

1 FTU is equivalent to the amount applied to two adult palms. Use this amount as a guide to determine the appropriate amount depending on the size of the affected area.



1 FTU(finger tip unit) Approximately 0.5g from the tip of the index finger to the first joint

Apply gently


Apply the cream on the affected area and gently spread it over the entire affected area without rubbing it in. The basic rule is to apply it until the surface becomes sticky.



Do not to rub it in

How to open the tube for the first time

Open the cap, turn it upside down, and insert the protrusion on the cap firmly into the opening of the tube to make a hole.



If symptoms do not improve or worsen after 5 to 6 days of use, stop using the medicine and consult with a doctor, pharmacist, or registered seller.

Side Effects

The frequency of side effects is low when topical steroids are used properly such as by the period of use and the amount applied. If used for a long period, the skin may become pale or thin. In addition, if used on areas infected with athlete's foot, the symptoms may worsen.

Attention for daily life

- ✓ The itching worsens when body gets warm, so bathe in lukewarm water for a short time.
- ✓ Wash body gently with a soft towel or your hands.
- ✓ Avoid clothing and bedding that come into direct contact with the skin and are made from materials such as wool or synthetic fibers.
- ✓ Avoid stimulants such as alcohol and spicy foods.

製品情報
Product features

