



For eczema, itching, and  
frostbite

Eczema, dermatitis treatment medicine

Product Name: Rinderon Vs Lotion

Class 2 Type Drug



# Rinderon Vs Lotion

This drug is a skin disease treatment, so do not use it under makeup or after shaving.  
Using large amounts or for a long period of time may cause side effects such as thinning of the skin and dilation of blood vessels in the skin. Take special care with the skin on the face, as it is thin.  
In addition, do not continue using the drug indiscriminately after symptoms have improved.



Please read  
carefully.



## Precautions



### When not to use this product

(If do not follow these instructions, current symptoms may worsen or adverse reactions are more likely to occur.)

- This product should not be used in the following persons.**  
Patients who have had an allergic symptom to this drug or its ingredients
- The medicine should not be used on the following parts:**
  - Varicella (chickenpox), athlete's foot/tinea or purulent lesions
  - Eyes and periorbital area
- Do not take extensively on the face.**
- Do not take this medicine for a long time.**



## Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.**  
(1) Patients undergoing medical treatment from a doctor (2) Pregnant women or women suspected of being pregnant (3) Patients who have experienced allergic symptoms associated with drugs, etc. (4) Those with extensive lesions (5) Those with severe moistening or erosion
- If the following symptoms are observed after using this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.**
- When symptoms do not improve even after using this drug medicine 5 to 6 times, stop using the drug and consult a physician, pharmacist or registered salesperson, being sure to take this document with you.**

Affected area	Symptoms
Skin	Rash/redness, itching
Skin (affected lesion)	Dermatomycosis such as athlete's foot/tinea, acne, purulent symptoms, persistent irritation, whitening



## Indications

Eczema, dermatitis, miliaria, irritated skin, itching, chilblain, frostbite, urticaria



## Dosage and Administration

Apply an appropriate amount to the affected lesion once or several times a day.

### <Precautions for dosage and Administration>

- Strictly follow the prescribed dosage and administration.
- When allowing children to use the medicine, a guardian must be present to watch and tell such them how to take it.
- Attention should be paid so that the medicine will not come into contact with eyes. If the medicine comes into contact with eyes, immediately rinse them with water or lukewarm water. If the symptoms are severe, seek an ophthalmologist for medical treatment.
- Shake well before using this medicine
- This medicine is for external use only.
- Do not cover the area with something that does not breathe, such as plastic wrap. When using on areas that come into contact with a diaper, do not use with tight-fitting diapers or sealed pants such as vinyl.
- Do not use under makeup or after shaving.



## Contact Desk

For inquiries about this product, please contact the store where you purchased it or contact the following:

●Shionogi Healthcare Co., Ltd. "Drug Information Center"  
Tel: Osaka 06-6209-6948, Tokyo 03-3406-8450  
Reception hours: 9:00-17:00 (excluding Saturdays, Sundays and holidays)



## Ingredient and amounts

Rinderon Vs Lotion is a white lotion, containing the following amount of ingredient per 1 gram:

Ingredient	Per 1g	Mode of action
Betamethasone valerate	1.2mg	A corticosteroid to suppress inflammation and to relieve itching

### Additives:

White petroleum jelly, liquid paraffin, cetanol, polyoxyethylene stearyl ether, butyl parahydroxybenzoate, methyl parahydroxybenzoate, sodium dihydrogen phosphate, phosphoric acid, and sodium hydroxide



## Precautions for storage and handling

- Store in a cool, low humidity place away from direct sunlight.
- Keep out of reach of children.
- Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the product.)
- Do not use products after its expiration date.



For information on the "Relief System for Sufferers from Adverse Drug Reactions"

●Pharmaceuticals and Medical Devices Agency (PMDA)  
[https://www.pmda.go.jp/kenkouhigai\\_camp/index.html](https://www.pmda.go.jp/kenkouhigai_camp/index.html)  
Tel: 0120-149-931 (toll-free number)

# Rinderon Vs Lotion

Topical steroids are effective for skin problems in hairy areas such as the scalp and armpits. Use Rinderon Vs Lotion properly, which is easy to use on hairy areas, to help speed up the healing process.

Skin problems such as eczema, dermatitis, rashes and associated itching in hairy areas such as the scalp, armpits and delicate areas. If the unpleasant symptoms accompanied by itching persist for a long time, you may continue to scratch the affected area, which may worsen the condition or leave scars. The key to treating skin problems in hairy areas is to suppress inflammation early and improve symptoms quickly. Therefore, it is important to use topical steroids correctly. Rinderon Vs Lotion is a milky lotion-type topical steroid that is easy to use on hairy areas.

Effective for various skin problems, including hairy areas.



For other symptoms

● Eczema on the back, stomach, and arms that is difficult to treat with sticky ointment.

● Eczema on the body during sweating seasons such as summer

※Do not use on mucous membranes.

※Do not use if the affected area is infected and oozing.

Point

HOW TO USE

## Usage

The amount that can be applied to an area the size of two palms is roughly the amount of medicine placed on an adult's index finger from the tip to the first joint (approximately 0.5g).※Adjust the amount according to the area of the affected area.



## Clean the affected area and apply

Apply to the clean affected area after bathing and washing hair. When applying to the scalp, it is easier to apply if you thoroughly wipe off the moisture with a towel and leave it slightly damp rather than drying it completely with a hair dryer.



## Take Lotion in hands and apply it directly to the affected area with fingers.

Take an appropriate amount in your hand and apply it directly to the affected area.

Apply it to scalp, try to avoid getting it on hair as much as possible by brushing it back. The basic rule is to make sure the surface is moist.



## Caution

● Do not place the mouth of the container directly on the affected area, as this may contaminate the container and worsen the inflammation.

● For sensitive areas, use only on areas smaller than diameter 3cm. If the affected area is large, stop using the product and consult a doctor, pharmacist, or registered distributor.



If symptoms do not improve or worsen after 5 to 6 days of use, stop using the medicine and consult a doctor, pharmacist, or registered seller.

## Use for skin symptoms associated with suppuration

Dermatitis often causes severe itching, and continued scratching weakens the barrier function that protects the skin from infection, making it easier for bacteria to enter through the skin. This medication improves symptoms such as dermatitis and itching. Do not use on areas with suppuration.

Condition	Dermatitis without Suppuration (Infection)	Dermatitis with Suppuration (Infection)
Use or Don't	○	×



Don't use as eye drops

※Long-term use around the eyes may increase intraocular pressure, so discontinue use.

## Side Effects

The frequency of side effects is low when topical steroids are used properly such as by the period of use and the amount applied. If used for a long period, the skin may become pale or thin. In addition, if used on areas infected with athlete's foot, the symptoms may worsen.



製品特徴  
Product features