



For eczema, itching, and frostbite

Eczema, dermatitis treatment medicine

Product name: Rinderon Vs Premium Ointment

Rinderon Vs Premium Ointment



This drug is a skin disease treatment, so do not use it under makeup or after shaving. Using large amounts or for a long period of time may cause side effects such as thinning of the skin and dilation of blood vessels in the skin. Take special care with the skin on the face, as it is thin. In addition, do not continue using the drug indiscriminately after symptoms have improved.



Please read carefully.

⚠ Precautions



When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

- This product should not be used in the following persons.**
Patients who have had an allergic symptom to this drug or its ingredients
- The medicine should not be used on the following parts:**
 - Varicella (chickenpox), athlete's foot/tinea or purulent lesions
 - Eyes and periorbital area

- Do not take extensively on the face.**
- Do not take this medicine for a long time.**



Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.**
 - Patients undergoing medical treatment from a physician
 - Pregnant women or women suspected of being pregnant
 - Patients who have experienced allergic symptoms associated with drugs, etc.
 - Those with extensive lesions
 - Those with severe moistening or erosion
- If the following symptoms are observed after using this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.**
- When symptoms do not improve even after using this drug 5 to 6 times, stop using the drug and consult a physician, pharmacist or registered salesperson, being sure to take this document with you.**

Affected area	Symptom
Skin	Rash/redness, itching
Skin (affected lesion)	Dermatomyces such as athlete's foot/tinea, acne, purulent symptoms, persistent irritation, whitening



Indications

Eczema, dermatitis, frostbite, miliaria, irritated skin, itching, chilblain, urticaria



Dosage and administration

Apply an appropriate amount to the affected lesion once to several times a day

< Precautions for Dosage and Administration >

- Strictly follow the prescribed dosage and administration.
- When allowing children to use the medicine, a guardian must be present to watch and tell such them how to use it.
- Attention should be paid so that the medicine will not come into contact with eyes. If the medicine comes into contact with eyes, immediately rinse them with water or lukewarm water. If the symptoms are severe, seek an ophthalmologist for medical treatment.
- This medicine is for external use only.
- Do not cover the area with something that does not breathe, such as plastic wrap. When using on areas that come into contact with a diaper, do not use with tight-fitting diapers or sealed pants such as vinyl.
- Do not use under makeup or after shaving.



Contact information

For inquiries about this product, please contact the store where you purchased it or the following:
• Shionogi Healthcare Co., Ltd. "Drug Information Center"
Tel: Osaka 06-6209-6948, Tokyo 03-3406-8450
Reception hours: 9:00-17:00 (excluding Saturdays, Sundays, and holidays)



Ingredients and amounts

Rinderon Vs Premium Ointment is white to pale yellow ointment, containing the following amounts of ingredients per 1 g of the ointment:

Ingredient	Amount (per 1 g)	Modes of action
Betamethasone valerate	1.2 mg	A corticosteroid to suppress inflammation and to relieve itching
Allantoin	2 mg	To help repair damaged skin tissue
Tocopherol acetate	10 mg	To improve blood flow and to prevent inflammation from spreading
Isopropyl methylphenol	1 mg	To exhibit a bactericidal effect on the skin, thereby preventing exacerbation due to infection

Additives: Liquid paraffin, glyceryl triisooctanoate, sorbitan sesquioleate, and white petrolatum.



Precautions for storage and handling

- Store in a cool, low humidity place away from direct sunlight.
- Keep out of reach of children.
- Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the product.)
- Do not use the product after its expiration date.

