



It supports display and voice reading in multiple languages.

Product features
Product features
QR Translator



This is the English version of the package insert of this product, translated from the Japanese package insert. The English version is provided for consumers who have purchased and use this product in Japan to help their understanding of its details. Please carefully read and fully understand the information given on this package insert before taking this product.

3 __ [underlined] parts: revised in January 2023

Cold remedy
(Non-pyrine)

Pylon PL Granules Pro

Product name: Pylon PL Granules Pro

Class ☒ OTC drug



Pylon PL Granules Pro is a non-pyrine cold medicine that has an excellent effect on various cold symptoms such as "sore throat," "fever," and "running nose" with the action of four active ingredients: Salicylamide and Acetaminophen, which are antipyretic and analgesic components; Promethazine methylenedisalicylate, which is an antihistamine component; and Anhydrous caffeine, which helps to suppress pain.



Please read carefully.



Precautions



When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

- This product should not be taken in the following persons.**
 - (1) Patients who have had an allergic symptom to this drug or its ingredients
 - (2) Patients who have experienced asthma from taking this drug or other cold remedies, or antipyretic analgesics
 - (3) Persons diagnosed as having the following: urination difficulty, gastric/duodenal ulcer, glaucoma
 - (4) Children under 15 years old
- This drug should not be taken together with the following drugs:**
Other cold medicines, antipyretics/analgesics, sedatives, expectorants, internal medicine, etc. containing antihistamines (internal medicine for rhinitis, medicine for motion sickness, medicine for allergies, etc.)
- After taking this drug, do not drive a car or operate machinery (sleepiness may occur).**
- Do not drink alcohol before/after taking this drug.**
- Do not take this drug for a long time.**



Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.**
 - (1) Patients undergoing medical treatment from a physician or dentist
 - (2) Nursing women, pregnant women or women suspected of being pregnant
 - (3) The elderly
 - (4) Patients who have experienced allergic symptoms associated with drugs, etc.
 - (5) Persons with the following symptom: hyperthermia
 - (6) Persons diagnosed as having the following: heart disease, liver disease, kidney disease, bronchial asthma
- If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.**

Affected area	Symptoms
Skin	Rash/redness, itching, oedema
Blood	Skin and mucous membranes become dark purple, anemia
Gastrointestinal system	Nausea and vomiting, loss of appetite, heartburn, abdominal pain, gastrointestinal bleeding
Neuropsychiatric system	Dizziness, malaise, headache, ear ringing, deafness, vision disorders, anxiety, hypersensitivity, insomnia, convulsions, mild clouding of consciousness, excitement, hallucination, delusion
Respiratory system	Cough, breathing difficulties
Cardiovascular system	Hypertension, hypotension, tachycardia
Urinary system	Urination difficulty, urinary retention
Other	Excessive temperature decrease, sweating, shivering, rhinitis symptoms, conjunctivitis

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid:

Name of symptom	Symptom
Shock (anaphylaxis)	Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take.
Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, acute generalised exanthematous pustulosis	Hyperthermia, ocular hyperaemia, eye discharge, lip erosion, pain throat, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen.
Drug-induced hypersensitivity syndrome	Widespread skin redness, systemic rash, fever, general malaise, swollen lymph nodes (in the neck, armpit, groin, etc.) may occur.
Aplastic anemia	Symptoms, such as bruises, nose bleeds, bleeding gums, fever, pale appearance of skin and mucosa, fatigue, palpitations, shortness of breath, becoming dizzy from feeling sick and blood in the urine, etc. may occur.
Agranulocytosis	Symptoms, such as acute high fever, chills and sore throat, etc. may occur.
Thrombocytopenia	Symptoms, such as nosebleeds, bleeding from the gums and bruises, caused by reduced counts of platelets which is a component of the blood may occur.
Asthma	Symptoms, such as wheezing or hissing when breathing, and difficult breathing, etc. may occur.
Interstitial pneumonia	Shortness and/or difficulties of breath when go upstairs or overwork, sudden dry cough and/or fever and its continuance
Hepatic function failure	Symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, general malaise, loss of appetite, etc. may occur.
Kidney disorders	Symptoms, such as fever, rash, reduced urinary volume, general oedema, general malaise, arthralgia (painful joints), diarrhea, etc. may occur.
Rhabdomyolysis	Symptoms may include pain in the muscles of the limbs/shoulders/lower back, numbness, weakness or stiffness in the limbs, general malaise and reddish-brown urine.
Glaucoma attack	Sudden ocular hyperaemia (congestion of the eye), eye pain, blurred vision, headache, and nausea may occur.

- The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation:**
Dry mouth, drowsiness
- When symptoms do not improve even after taking this drug 5 to 6 times, stop taking the drug and consult a physician, pharmacist or registered salesperson, being sure to take this document with you.**



Indication

Relief of various symptoms of a common cold: sore throat, fever, running nose, stuffy nose, sneezing, chills (feeling cold due to fever), headache, joint pain, and muscle pain.



Dosage and administration

Adults (15 years old and over): Take 1 sachet 4 times a day after each meal and before bed in intervals of at least 4 hours between doses, as a standard.

Age	Single dose	Daily dosing frequency
Adults (15 years old and over)	1 sachet	4 times a day
Children under 15 years old	Not allowed to take this medicine	

- Strictly follow the prescribed dosage and administration.



Ingredients and amounts

Pylon PL Granules Pro are white granules, containing the following amounts of ingredients per 4 sachets (1 g x 4 = the daily dose for adults):

Ingredient	Amount (per 4 sachets)	Modes of action
Salicylamide	1080 mg	To relieve pain and to reduce fever
Acetaminophen	600 mg	
Anhydrous caffeine	240 mg	To help relieve pain
Promethazine methylenedisalicylate	54 mg	To relieve running nose, stuffy nose, and sneezing

Additives include lactose hydrate, corn starch, sodium chloride, sucrose, and hydrated silicon dioxide.



Precautions for storage and handling

- Store in a cool, low humidity place away from direct sunlight.
- Keep out of reach of children.
- Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the drug.)
- Do not take the product after its expiration date.



Manufactured and Distributed by:
Shionogi Healthcare Co., Ltd.
2-6-18 Kitahama, Chuo-ku, Osaka