



Product information  
Product features



It supports display and  
voice reading in multiple  
languages. Scan the QR  
code.

This is the English version of the package insert of this product, translated from the Japanese package insert. The English version is provided for consumers who have purchased and use this product in Japan to help their understanding of its details. Please carefully read and fully understand the information given on this package insert before taking this product.

3

Antitussive drug (non-narcotic)

Product name: Medicon Cough Suppressant Tablets Pro

Class 2 OTC drug

# Medicon Cough Suppressant Tablets Pro

SHIONOGI

Medicon Cough Suppressant Tablets Pro are a non-narcotic antitussive drug that acts directly on the cough center, exerting an excellent effect against the distressing symptom of cough.



## Precautions



### When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

1. This product should not be taken in the following persons.  
Patients who have had an allergic symptom to this drug or its ingredients
2. This drug should not be taken together with the following drugs:  
Other antitussives and expectorants, cold medicines, sedatives, internal medicines, etc. containing antihistamines (internal medicine for rhinitis, medicine for motion sickness, medicine for allergies, etc.)
3. After taking this drug, do not drive a car or operate machinery (sleepiness may occur).



### Consultation

1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.
  - (1) Patients undergoing medical treatment from a physician
  - (2) Pregnant women or women suspected of being pregnant
  - (3) Patients who have experienced allergic symptoms associated with drugs, etc.
  - (4) Persons with the following symptom: hyperthermia
2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Affected area	Symptom
Skin	Rash/redness, itching
Gastrointestinal system	Nausea and vomiting, loss of appetite
Neuropsychiatric system	Dizziness
Respiratory system	Breathing difficulties, shortness of breath

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid:

Name of symptom	Symptom
Shock (anaphylaxis)	Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take.

3. The following symptom may be observed after taking this drug. If this symptom persists or worsens, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation:  
Drowsiness
4. When symptoms do not improve even after taking this drug 5 to 6 times, stop taking the drug and consult a physician, pharmacist or registered salesperson, being sure to take this document with you.



## Indications

Cough



## Dosage and administration

Take the following amount with water or lukewarm water.  
Take this medicine in intervals of at least 4 hours.

Age	Single dose	Daily dosing frequency
Adults (15 years old and over)	2 tablets	3 times a day
Children under 15 years old	Not allowed to take this medicine	

- Strictly follow the prescribed dosage and administration.

- How to take out the tablet

As shown in the right figure, press the raised part of the PTP sheet containing the tablet firmly with your fingertips to break the aluminum foil on the back, take out the tablet and swallow it. (If the PTP sheet is swallowed by mistake, it may lead to unexpected accidents such as piercing the esophageal mucosa.)

How to take out the tablet



## Ingredients and amounts

Medicon Cough Suppressant Tablets Pro are white tablets, containing the following amounts of ingredients per 6 tablets (the daily dose for adults):

Ingredient	Amount (per 6 tablets)	Mode of action
Dextromethorphan hydrobromide hydrate	90 mg	To relieve cough

Additives include lactose hydrate, corn starch, microcrystalline cellulose, magnesium stearate, talc, white shellac, and castor oil.



## Precautions for storage and handling

- (1) Store in a cool, low humidity place away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not remove from the PTP sheet and transfer to another container.  
(This may lead to misuse or alter the quality of the drug.)
- (4) Do not take the product after its expiration date.



## Contact information

For inquiries about this product, please contact the store where you purchased it or the following:

• Shionogi Healthcare Co., Ltd. "Drug Information Center"  
Tel: Osaka 06-6209-6948, Tokyo 03-3406-8450  
Reception hours: 9:00-17:00 (excluding Saturdays, Sundays, and holidays)

For information on the "Relief System for Sufferers from Adverse Drug Reactions"

• Pharmaceuticals and Medical Devices Agency (PMDA)  
[https://www.pmda.go.jp/kenkouhigai\\_camp/index.html](https://www.pmda.go.jp/kenkouhigai_camp/index.html)  
Tel: 0120-149-931 (toll-free number)

Manufactured and Distributed by:

**Shionogi Healthcare Co., Ltd.**  
2-6-18 Kitahama, Chuo-ku, Osaka