



It supports display and voice reading in multiple languages.

Product features
Product features
QR Translator



This is the English version of the package insert of this product, translated from the Japanese package insert. The English version is provided for consumers who have purchased and use this product in Japan to help their understanding of its details. Please carefully read and fully understand the information given on this package insert before taking this product.

1

Antitussive and expectorant drug (non-narcotic)

Product name: Medicon Cough Suppressant Syrup Pro

Class 2 OTC drug

Medicon Cough Suppressant Syrup Pro



Medicon Cough Suppressant Syrup Pro is a non-narcotic antitussive drug that exerts excellent effects against the distressing symptoms of **cough and sputum** by the actions of 2 active ingredients: dextromethorphan hydrobromide hydrate that acts directly on the cough center to suppress cough, and potassium cresolsulphonate that facilitates sputum expectoration.



Please read carefully.



Precautions



When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

1. This product should not be taken in the following persons.
Patients who have had an allergic symptom to this drug or its ingredients
2. This drug should not be taken together with the following drugs:
Other antitussives and expectorants, cold medicines, sedatives, internal medicines, etc. containing antihistamines (internal medicine for rhinitis, medicine for motion sickness, medicine for allergies, etc.)
3. After taking this drug, do not drive a car nor operate machinery (sleepiness may occur).



Consultation

1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.
 - (1) Patients undergoing medical treatment from a physician
 - (2) Pregnant women or women suspected of being pregnant
 - (3) Patients who have experienced allergic symptoms associated with drugs, etc.
 - (4) Persons with the following symptom: hyperthermia
2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Affected area	Symptom
Skin	Rash/redness, itching
Gastrointestinal system	Nausea and vomiting, loss of appetite
Neuropsychiatric system	Dizziness
Respiratory system	Breathing difficulties, shortness of breath

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid:

Name of symptom	Symptom
(anaphylaxis)	Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take.

3. The following symptom may be observed after taking this drug. If this symptom persists or worsens, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation: Drowsiness
4. When symptoms do not improve even after taking this drug 5 to 6 times, stop taking the drug and consult a physician, pharmacist or registered salesperson, being sure to take this document with you.
5. If this medicine is taken for more than 2 weeks even after symptoms have been improved, consult a physician, pharmacist, or registered salesperson for advice.



Indications

Cough, sputum



Dosage and administration

Take the following dose amount 3 times a day. Take this medicine in intervals of at least 4 hours.


- Strictly follow the prescribed dosage and administration.

Age	Single dose
Adults (15 years old and over)	6 mL
Children under 15 years old	Not allowed to take this medicine

A safety cap that cannot be easily opened by children is used for the container.

How to use the safety cap

Open the cap by turning it counterclockwise while pushing it downward.



Correct oral intake method

Place the mouth of the bottle onto the edge of the measuring cup and pour a single dose amount (6 mL) of the drug solution into the cup. Only the specified dose amount must be taken at one time.



* Stand and hold the bottle on a flat surface.



Ingredients and amounts

Medicon Cough Suppressant Solution Pro is a clear, pale yellowish-brown, viscous solution, containing the following amounts of ingredients per 18 mL of the solution (the daily dose for adults):

Ingredient	Amount (per 18 mL)	Modes of action
Dextromethorphan hydrobromide hydrate	45 mg	To relieve cough
Potassium cresolsulphonate	270 mg	To facilitate sputum expectoration

Additives include ethanol, sucrose, caramel, benzoic acid, sodium hydroxide, benzyl benzoate, ethyl vanillin, glycerin, vanillin, and flavoring.



Precautions for storage and handling

- (1) Store in a cool, low humidity place away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the drug.)
- (4) Wash the measuring cup with water after each use and keep it clean.
- (5) When closing the cap, thoroughly wipe the mouth of the bottle. If the cap is closed with some drug solution attached to the mouth of the bottle, this may cause the cap not to be easily opened at the next use. If the cap is not tightly closed, this may cause leakage or contamination of the solution from the bottle.
- (6) If the cap is not easily opened, turn the bottle over and immerse the cap in hot water (50°C to 60°C) to enable opening easily.
- (7) Do not take the product after its expiration date.



Contact information

For inquiries about this product, please contact the store where you purchased it or the following:

- Shionogi Healthcare Co., Ltd. "Drug Information Center"

Tel: Osaka 06-6209-6948, Tokyo 03-3406-8450

Reception hours: 9:00-17:00 (excluding Saturdays, Sundays, and holidays)

For information on the "Relief System for Sufferers from Adverse Drug Reactions"

- Pharmaceuticals and Medical Devices Agency (PMDA)

https://www.pmda.go.jp/kenkouhigai_camp/index.html

Tel: 0120-149-931 (toll-free number)

Manufactured and Distributed by:

Shionogi Healthcare Co., Ltd.

2-6-18 Kitahama, Chuo-ku, Osaka